

Data Sheet

Product Name:	Avelumab
Cat. No.:	CS-0030892
CAS No.:	1537032-82-8
Target:	PD-1/PD-L1
Pathway:	Immunology/Inflammation
Solubility:	H ₂ O

Avelumab

BIOLOGICAL ACTIVITY:

Avelumab is a fully human IgG1 anti-**PD-L1** monoclonal antibody with potential antibody-dependent cell-mediated cytotoxicity. IC₅₀ & Target:PD-1/PD-L1^[1] *In Vitro*: Avelumab is a fully human IgG1 anti-PD-L1 monoclonal antibody with potential antibody-dependent cell-mediated cytotoxicity property. Avelumab increases NK-cell lysis 3.1-fold (P=0.01) in JHC7 cells relative to isotype control. When the cells are treated with IFN- γ , Avelumab markedly enhances NK-cell lysis relative to isotype control in the following cell lines: JHC7 (7.56-fold; P=0.001), UM-Chor1 (7.34-fold; P<0.001), U-CH2 (2.6 fold; P=0.008), MUG-Chor1 (8.38-fold; P=0.0016). Avelumab effectively increases antibody-dependent cell-mediated cytotoxicity (ADCC) of both the non-cancer stem cell (CSC) and CSC subpopulations to the same degree^[1]. Results also demonstrate that the addition of Avelumab increases the frequency of antigen-specific multifunctional CD8⁺ T cells by more than fivefold, relative to the isotype control in CEFT-stimulated peripheral blood mononuclear cells (PBMCs)^[2]. *In Vivo*: Measurement of individual tumors clearly shows a slowing of tumor growth in the Avelumab-treated mice. By day 36 post-tumor implantation, there is a significant (P<0.01) reduction in the average tumor volume of the Avelumab-treated mice. Reduction in MB49 tumor growth in the mice treated with Avelumab is durable and leads to a significant (P<0.05) improvement in percent survival. Avelumab treatment of 10 mice with bladder tumors results in complete tumor regression in 8 mice, confirmed by histopathology. However, in mice depleted of either CD4 or CD8 cells, Avelumab treatment is much less effective in controlling bladder tumor burden with tumor breakthrough occurring in a higher frequency in mice depleted of CD4 T cells^[3].

PROTOCOL (Extracted from published papers and Only for reference)

Cell Assay: ^[1]To examine the relationship between a cancer stem cell (CSC) subpopulation and antibody-dependent cell-mediated cytotoxicity (ADCC) activity, **UM-Chor1 cells** are left untreated or treated with 50 ng/mL of IFN- γ for 24 h. Cells are then plated as targets at 50,000 cells/well in 6-well round-bottom culture plates and incubated with **2 μ g/mL of Avelumab** at room temperature for **30 min**. NK cells are added at 2500,000 cells/well at an effector-to-target (E:T) ratio of 50:1. After 4 h, tumor cells are harvested and stained with antibodies for flow cytometry^[1]. **Animal Administration:** ^[3]**Female C57BL/6 mice** are used in this study. Subcutaneous tumor injections are carried out by inoculating C57BL/6 mice with 1 \times 10⁵ MB49 parental cells on the right shaved flank. Tumor growth is measured with calipers and 8 days post-inoculation mice are assigned to treatment groups. Tumor-bearing mice are treated with **Avelumab (400 μ g per 100 μ L)** and injected i.p. **three times, 3 days apart**. Since Avelumab is a human IgG1, three injections have to be compressed within a **7 to 9 day** window (i.e., days 9, 12, and 15 post-tumor inoculation) to avoid the onset of neutralizing mouse anti-human Ig^[3].

References:

- [1]. Fujii R, et al. Enhanced killing of chordoma cells by antibody-dependent cell-mediated cytotoxicity employing the novel anti-PD-L1 antibody avelumab. *Oncotarget*. 2016 Jun 7;7(23):33498-511.
- [2]. Grenga I, et al. A fully human IgG1 anti-PD-L1 MAb in an in vitro assay enhances antigen-specific T-cell responses. *Clin Transl Immunology*. 2016 May 20;5(5):e83.
- [3]. Vandeveer AJ, et al. Systemic Immunotherapy of Non-Muscle Invasive Mouse Bladder Cancer with Avelumab, an Anti-PD-L1 Immune Checkpoint Inhibitor. *Cancer Immunol Res*. 2016 May;4(5):452-62.

CAIndexNames:

Immunoglobulin G1, anti-(human CD antigen CD274) (human monoclonal MSB0010718C heavy chain), disulfide with human monoclonal MSB0010718C light chain, dimer

SMILES:

[Avelumab]

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 610-426-3128

Fax: 888-484-5008

E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite F, Monmouth Junction, NJ 08852, USA