


Data Sheet

Product Name:	GENE-6776
Cat. No.:	CS-0031103
CAS No.:	2009273-71-4
Molecular Formula:	C ₂₀ H ₂₀ N ₄ O ₂
Molecular Weight:	348.40
Target:	Deubiquitinase
Pathway:	Cell Cycle/DNA Damage
Solubility:	DMSO : ≥ 100 mg/mL (287.03 mM)



BIOLOGICAL ACTIVITY:

GENE-6776 is a selective and orally bioavailable **USP7** inhibitor^[1]. IC₅₀ & Target: USP7^[1] *In Vitro*: GNE-6776 significantly inhibits USP7 at 15 μM. GNE-6776 is a highly selective USP7 inhibitor against both recombinant and endogenous cellular deubiquitinases^[1]. *In Vivo*: GNE-6776 (100 or 200 mg/kg; oral gavage on a once or twice daily schedule; for 10 days) inhibits EOL-1 xenograft growth in mice^[1].

PROTOCOL (Extracted from published papers and Only for reference)

Cell Assay: ^[1]EOL-1 cells are seeded into 384-well plates 24 h before compound addition. Cells are then incubated with compound (e.g., **GENE-6776; 0.003, 0.009, 0.027, 0.082, 0.25, 0.74, 2.22, 6.67, and 20 μM**) for 72 h or 120 h before assaying viability. Assays are performed in biological triplicate. Cells are incubated (37°C, 5% CO₂) in RPMI-1640, 2.5% FBS (72 h assay) or 5% FBS (120 h assay), and 2 mM glutamine throughout the assay. The reported IC₅₀ and mean viability metrics are as follows: IC₅₀ is the dose at which the estimated inhibition is 50% relative to untreated wells (that is, absolute IC₅₀). The mean viability is calculated^[1].

Animal Administration: GNE-6776 is formulated as a suspension in 0.5% methylcellulose/0.2% Tween-80^[1].^[1]Mice^[1]

GENE-6776 is administered at **200 mg/kg** (body weight) by **oral gavage** to **female C.B-17 SCID mice**, aged 12-16 weeks (n=3 per time point). No randomization is used for DMPK studies. At 0.5, 1, 2, 4, 8 and 24 h post-dose, blood samples are collected by terminal cardiac puncture into anticoagulant tubes (EDTA). Clarified plasma is then transferred to a fresh tube and snap frozen. GNE-6776 plasma concentrations are determined by LC-MS/MS^[1].

References:

[1]. Lorna Kategaya , et al. USP7 Small-Molecule Inhibitors Interfere With Ubiquitin Binding. Nature. 2017 Oct 26;550(7677):534-538.

CAIndexNames:

[3,3'-Bipyridine]-6-carboxamide, 6'-amino-4'-ethyl-5'-(4-hydroxyphenyl)-N-methyl-

SMILES:

CCC(C(C1=CC=C(C(O)C=C1)=C(N)N=C2)=C2C3=CC=C(C(C(NC)=O)N=C3

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 610-426-3128

Fax: 888-484-5008

E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA