

Data Sheet

Product Name:	Chitosan oligosaccharide	
Cat. No.:	CS-0043357	
CAS No.:	148411-57-8	
Target:	AMPK; Endogenous Metabolite	
Pathway:	Epigenetics; Metabolic Enzyme/Protease; PI3K/Akt/mTOR	Chitosan oligosaccharide
Solubility:	H ₂ O : 50 mg/mL (Need ultrasonic); DMSO : 50 mg/mL (Need ultrasonic)	

BIOLOGICAL ACTIVITY:

Chitosan oligosaccharide (COS) is an oligomer of β -(1 $\rightarrow$ 4)-linked D-glucosamine. Chitosan oligosaccharide (COS) activates **AMPK** and inhibits inflammatory signaling pathways including **NF- κ B** and **MAPK** pathways. IC₅₀ & Target: AMPK^[1] **In Vitro:** Chitosan oligosaccharide (COS) represents a class of natural polymers that holds therapeutic promise in several diseases due to not only its physiochemical properties including water-solubility and low viscosity but also its favorable pharmacological properties including good pharmacokinetics and safety profiles and diverse beneficial biological activities. Activation of AMPK and inhibition of inflammatory signaling pathways including NF- κ B and MAPK pathways are recognized as major mechanisms responsible for several effects of Chitosan oligosaccharide (COS) including anti-inflammation, anti-cancer, and anti-diabetes. COS can interrupt cancer progression at multiple stages by modulating several signaling proteins/pathways including NF- κ B, AMPK, mTOR, caspase-3, CD147, MMP-2, MMP-9, and VEGF. In vitro experiments have demonstrated that Chitosan oligosaccharide (COS) induced the death of several cancer cell types including ascites, bladder cancer, prostate cancer, lung cancer, liver cancer, leukemia, cervical cancer and colorectal cancer. The values of IC₅₀ of Chitosan oligosaccharide (COS) in inducing cytotoxicity are 25 μ g/mL-50 μ g/mL depending on types of cancer cells^[1]. **In Vivo:** The oral administration of Chitosan oligosaccharide (16 mg/kg/day) suppresses the production of the proinflammatory cytokines involved in allergic reactions, i.e., IL-4, IL-13 and TNF- α , in the lung tissues and bronchoalveolar lavage fluid of the mice. Last, an anti-inflammatory effect of Chitosan oligosaccharide (COS) on lymphocyte activation has been documented in a rat model of autoimmune anterior uveitis induced by immunization with bovine melanin-associated antigen^[1]. Chitosan oligosaccharide (COS) inhibits UV-induced macroscopic appearance in mice skin. Compared with healthy dorsal skin with smoothness and some shallow wrinkles of hairless mice in normal control group, UV exposure for 10 weeks triggers skin erythema, dry, thickening, sagging and coarse wrinkles, and even leathery appearance and slight flesh-colored lesion in the model mice, the visual score of which is markedly higher than that of the normal control group ($p < 0.05$), indicating that UV induces photoaging in skin surface^[2].

PROTOCOL (Extracted from published papers and Only for reference)

Animal Administration: ^[2]Mice^[2]

Seven-week-old female hairless BALB/c mice (n=12), weighing approximately 16 g, are used. After a week of acclimation, the hairless mice are randomly divided into five groups of 6 mice per group: Normal control group (without UV irradiation but with double-distilled water treatment); Model group (UV irradiation with double-distilled water treatment); COS-L group, COS-M group and COS-H group (UV irradiation with **50 mg/mL**, **100 mg/mL** and **200 mg/mL COS** treatment, respectively). For mice used for topical application, 100 μ L are applied to each mouse dorsal skin every time after UV-radiation. The dorsal treated skin area of mice is carefully wiped with soft absorbent cotton soaked in distilled water, and then wiped with dry cotton before each UV-radiation to

remove any remaining Chitosan oligosaccharide (COS)^[2].

References:

[1]. Muanprasat C, et al. Chitosan oligosaccharide: Biological activities and potential therapeutic applications. Pharmacol Ther. 2017 Feb;170:80-97

[2]. Kong SZ, et al. Anti-photoaging effects of chitosan oligosaccharide in ultraviolet-irradiated hairless mouse skin. Exp Gerontol. 2018 Mar;103:27-34.

CAIndexNames:

Chitosan, 2-hydroxypropanoate (ester)

SMILES:

[Chitosan oligosaccharide]

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 610-426-3128

Fax: 888-484-5008

E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA