


Data Sheet

Product Name:	3-(Methylthio)propionic acid
Cat. No.:	CS-6296
CAS No.:	646-01-5
Molecular Formula:	C ₄ H ₈ O ₂ S
Molecular Weight:	120.17
Target:	Endogenous Metabolite; Fungal
Pathway:	Anti-infection; Metabolic Enzyme/Protease
Solubility:	H ₂ O : 25 mg/mL (208.04 mM; Need ultrasonic)



BIOLOGICAL ACTIVITY:

3-(Methylthio)propionic acid is an intermediate in the methionine metabolism. **In Vitro:** Methionine has consistently been shown to be the most toxic amino acid in experiments devised to assess the relative toxicity of dietary amino acids. 3-methylthiopropionate is an intermediate in methionine catabolism in rat and monkey liver in vitro. This pathway appears to account for a major portion of methionine oxidation in vitro^[1]. Cultures of *Streptomyces lincolnensis* accumulated 3-methylthioacrylic acid in amounts directly related to the concentration of methionine in the medium. The first intermediate in the pathway may be the keto acid, which is then oxidatively decarboxylated to 3-methylthiopropionic acid^[2]. The purified 3-MTPA has antifungal activity in assays using *F. oxysporum* as a model fungus. Daily measurements of shoot and root length shows severe inhibition of seed germination and root and shoot development at concentrations above 12mg for partially purified extracts^[3]. 3-methylthiopropionic acid ethyl ester possesses potential anticarcinogenic properties by inducing differentiation in well-differentiated colon cancer cells. Treatment of RCM-1 cells for 4 days with 3-methylthiopropionic acid ethyl ester between the doses of 0.25 and 2 mM progressively increases the percent area occupied by duct structures relative to the control, and also induces an increase in the number and the maximum diameter of the ducts in each culture plate^[4].

PROTOCOL (Extracted from published papers and Only for reference)

Cell Assay: ^[4]RCM-1 cells are each plated into 35-mm plastic culture plates. Twenty-four hours after plating, cells are treated with 3-methylthiopropionic acid ethyl ester in 2 mL of 10% FBS-RPMI+F12 for 4 days, and are counted by using a hemacytometer, after trypsinization with 0.25% trypsin and 1 mM EDTA^[4].

References:

- [1]. Steele RD, et al. Identification of 3-methylthiopropionic acid as an intermediate in mammalian methionine metabolism in vitro. *J Biol Chem.* 1978 Nov 10;253(21):7844-50.
- [2]. Surette R, et al. Formation of 3-methylthioacrylic acid from methionine by *Streptomyces lincolnensis*. Isolation of a peroxidase. *J Antibiot (Tokyo).* 1976 Jun;29(6):646-52.
- [3]. Kim YC, et al. 3-methylthiopropionic acid produced by *Enterobacter intermedium* 60-2G inhibits fungal growth and weed seedling development. *J Antibiot (Tokyo).* 2003 Feb;56(2):177-80.

[4]. Nakamura Y, et al. 3-Methylthiopropionic acid ethyl ester, isolated from Katsura-uri (Japanese pickling melon, Cucumis melo var. conomon), enhanced differentiation in human colon cancer cells. J Agric Food Chem. 2008 May 14;56(9):2977-84.

CAIndexNames:

Propanoic acid, 3-(methylthio)-

SMILES:

O=C(O)CCSC

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 610-426-3128

Fax: 888-484-5008

E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA