


Data Sheet

Product Name: H-cis-Hyp-OMe (hydrochloride)

Target: ADC Linkers; PROTAC Linkers

Pathway: Antibody-drug Conjugate/ADC Related; PROTAC

Solubility: 10 mM in DMSO



HCI

BIOLOGICAL ACTIVITY:

H-cis-Hyp-OMe hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). H-cis-Hyp-OMe hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PR **In Vitro**: ADCs are comprised of an antibody to which is attached an ADC cytotoxin through an ADC linker^[1].

PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins^[2].

References:

- [1]. Beck A, et al. Strategies and challenges for the next generation of antibody-drug conjugates. Nat Rev Drug Discov. 2017;16(5):315-337.
- [2]. Nalawansha DA, et al. PROTACs: An Emerging Therapeutic Modality in Precision Medicine. Cell Chem Biol. 2020;27(8):998-985.

CAIndexNames:

L-Proline, 4-hydroxy-, methyl ester, hydrochloride (1:1), (4S)-

SMILES:

O=C([C@H]1NC[C@@H](O)C1)OC.CI

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 610-426-3128 Fax: 888-484-5008 E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr., Suite Q., Monmouth Junction, NJ 08852, USA

Page 1 of 1 www.ChemScene.com